arXiv:hep-th/0207008v1 1 Jul 2002

On the effects of the vanishing of the non-metricity to the effective string field equations

Abstract

Effective string field equations with zero-constrained torsion have been studied extensively in the literature. But, one may think that the effects of vanishing of the non-metricity have not been explained in detail and also in according to some recent literature [4],[5], the action density in my previous paper [3] is not complete. For these reasons, in this erratum, in according to the effects of vanishing of the non-metricity to the field equations, the action density of my previous paper [3] will be completed in a variational setting. Furthermore, up to now, an unambiguous derivation of the dilaton potential has not been given. If one thinks that vanishing of the non-metricity gives a spontaneous breakdown of local Weyl invariance then a dilaton potential is obtained unambiguously in this framework.

1.Introduction and Motivation

Effective string field equations with zero-constrained torsion have been studied extensively in the literature. But, one may think that the effects of vanishing of the non-metricity have not been explained in detail and also in according to some recent literature [4],[5], the action density in my previous paper [3] is not complete. For these reasons, in this erratum, in according to the effects of vanishing of the non-metricity to the field equations, the action density of my previous paper [3] will be completed in a variational setting. Furthermore, up to now, an unambiguous derivation of the dilaton potential has not been given. If one thinks that vanishing of the non-metricity gives a spontaneous breakdown of local Weyl invariance then a dilaton potential is obtained unambiguously in this framework.

2.Theory

In my previous paper [3], the first-order sector of the Lagrangian was

$$\begin{aligned}\mathcal{L} = e^{-\phi}(R_{ab}\wedge^{*} e^{ab} + Tr(F\wedge^{*} F) - \alpha d\phi\wedge^{*} d\phi + \beta H\wedge^{*} H)\\ +\gamma e^{-\frac{N-4}{N-2}\phi}R_{ab}\wedge R_{cd}\wedge^{*} e^{abcd}\\ +(dH - \epsilon R_{ab}\wedge R^{ab} - \epsilon' Tr(F\wedge F))\wedge\mu + T^{a}\wedge\lambda_{a}.\end{aligned} \tag{1}$$

To handle zero-constrained torsion effectively, some Lagrange multiplier $(N-2)-$ forms λ_a are introduced. But, unfortunately the vanishing of the torsion does not necessarily give the vanishing of the non-metricity, namely, from Eq.(2.9) in [4]

$$D^{*}e^{a} = \ ^{*}(e^{a} \wedge e_{b}) \wedge T^{b} + 4\, Q \wedge \ ^{*}e^{a}, \tag{2}$$

is satisfied, where Q is a non-metricity object. For the vanishing of the non-metricity, $f_a\ D^{*}e^{a}$ term will newly be added in the Eq.(1) action density, where f_a is a zero-form. The variation with respect to f_a gives $D^{*}e^{a} = 0$, so Q, namely, non-metricity equals to zero. The variation with respect to the ω has not been affected by this $f_a\ D^{*}e^{a}$ term because the ω variation term, $\omega^{ab} \wedge f_a\ ^{*}e_b$ equals identically to zero. So only the variation with respect to the e^a term rests. This term is

$$\delta e^{a} \wedge \ ^{*}(e^{c} \wedge e_{a}) \wedge D f_{c}. \tag{3}$$

There is no any other restriction on f_c. So, as a simple solution, if one selects $f_c = i_c dV(\phi)$ then the dilaton potential $V(\phi)$ is obtained.

In my previous paper [3], in the cosmological solutions, f_c has been taken as zero.

[1]. T.Dereli, R.W.Tucker,Class.Q.Grav.**4** (1987) 791
[2]. T.Dereli, Ş. Özkurt, Class.Q.Grav.**8** (1991) 889
[3]. Ş. Özkurt, Class.Q.Grav.**13** (1996) 265
[4]. T.Dereli, R.W.Tucker, JHEP 03 (2002) 041
[5]. D.Grumiller, W.Kummer, D.V.Vassilevich, hep-th 04 (2002) 253